

visited. We have thus established that there is a difference in the movement of membrane proteins in living and dead cells. The active component of the motion of the λ -receptor in living cells has been modeled as an artificial temperature, which estimates the energy necessary for this active motion. The influence of antimicrobial peptides (AMPs) on the outer membrane of bacteria was investigated using the mobility of the λ -receptor as a membrane marker. With the growing resistance to antibiotics AMPs are gaining increased interest. Using the AMPs polymyxin B (PMB), and the non-toxic derivative polymyxin B nonapeptide (PMBN), we have investigated the influence of AMPs on the outer bacterial membrane. Cells exposed to PMB showed a decrease in the spread of position visited by the λ -receptor upon poisoning. PMBN is known to increase the permeability of the outer membrane without killing the cells. Exposure to PMBN did, however, not influence the mobility of the λ -receptor.

1430-Pos Board B274

Investigating Axonal Outgrowth and Orientation of Neuroblasts through an Alternating Stiffness Substrate

Szu-Yuan Chou¹, Chao-Min Cheng¹, Maarten P. de Boer¹,
Chih-Cheng Chen², Philip R. LeDuc¹.

¹Carnegie Mellon University, Pittsburgh, PA, USA, ²Academia Sinica, Pittsburgh, Taiwan.

At the interface of cell-substrate interactions, substrate elasticities strongly influence the morphology and function of cellular responses. This is important in diverse areas including neural function, metastasis, and heart disease. Cells are subject to mechanical signals in addition to biochemical signals; therefore, understanding the cellular interactions to substrate stiffness and with extracellular matrix is an important step to define how cultured cells respond when grown on materials that have similar characteristics to physiological conditions. To address the effects of localized elasticity, we developed a new method to control the microenvironment through generating a substrate with localized alternating stiffnesses interacting with cells to affect their structural response. This technique was accomplished through first fabricating polymeric microchannels using conventional soft lithography. We made channels either 30 μm or 100 μm wide, and 50 μm deep with poly(dimethylsiloxane) (PDMS) with a 5:1 ratio of base/curing agent. We then poured PDMS with a 30:1 ratio of base/curing agent into the channels and then removed the extra PDMS to produce a level surface. This produced alternating surfaces with elastic modulus of 800 kPa and 200 kPa adjacent to one another. We coated the surfaces with extracellular matrix and seeded neuroblasts onto the systems. We then differentiated them using retinoic acid (20 μM). We found that the neuroblasts had distinct patterns that emerged as they extended processes to and across these alternating stiffness substrates. Depending on the location of the cell body and the direction of the outgrowth when compared to the alternating stiffness interface, the processes would extend forward in different paths. We believe that this approach will enable greater understanding of axon outgrowth as well as provide insight into a variety of diseases linked to cell-ECM-material interactions.

1431-Pos Board B275

FRAP and Photoconversion in Multiple Arbitrary Regions of Interest Using a Programmable Array Microscope (PAM)

Guy M. Hagen^{1,2}, Wouter Caarls¹, Keith A. Lidke^{1,3},
Anthony H.B. de Vries¹, Cornelia Fritsch^{1,4}, B. George Barisas⁵,
Donna J. Arndt-Jovin¹, Thomas M. Jovin¹.

¹Max Planck Institute for Biophysical Chemistry, Goettingen, Germany, ²1st Faculty of Medicine, Charles University in Prague, Prague, Czech Republic, ³University of New Mexico, Albuquerque, NM, USA, ⁴University of Sussex, Brighton, United Kingdom, ⁵Colorado State University, Fort Collins, CO, USA.

Photomanipulation (photobleaching, photoactivation, or photoconversion) is an essential tool in fluorescence microscopy. Fluorescence recovery after photobleaching (FRAP) is commonly used for the determination of lateral diffusion constants of membrane proteins, and can be conveniently implemented in confocal laser scanning microscopy (CLSM). Such determinations provide important information on molecular dynamics in live cells. However, the CLSM platform is inherently limited for FRAP because of its inflexible raster (spot) scanning format. We have implemented FRAP and photoconversion protocols using a programmable array microscope (PAM). The bleaching or photoconversion patterns are arbitrary in number and shape, dynamic, and adjustable to and by the sample characteristics. We have used multi-spot PAM-FRAP to measure the lateral diffusion of the erbB3 (HER3) receptor tyrosine kinase labeled by fusion with mCitrine on untreated cells and after treatment with reagents that perturb the cytoskeleton or plasma membrane or activate coexpressed erbB1 (HER1, the EGF receptor EGFR). We also explored the versatility of the PAM for photoconversion in arbitrary regions of interest, in cells expressing erbB3 fused with the fluorescent protein dropna.

1432-Pos Board B276

Glycosphingolipids and Non-Raft Phospholipids Exhibit Very Similar Dynamics in Single-Molecule Observations

Kenichi G.N. Suzuki¹, Takahiro K. Fujiwara², Akihiro Kusumi².

¹PRESTO-JST, iCeMS, Kyoto Univ., Kyoto, Japan, ²ICORP-JST, iCeMS, Kyoto Univ., Kyoto, Japan.

The diffusion of typical raft-associated molecules, glycosphingolipids, GM₁, GM₃, GD_{1a}, was observed using high-speed single-particle tracking (HS-SPT) at a temporal resolution of $\sim 6 \mu\text{s}$. Each glycosphingolipid molecule was tagged with a 40-nm-gold particle conjugated by the respective Fab antibody or a Cy3-conjugated Fab antibody. At video rate, both probes exhibited the same diffusion behavior, undergoing apparent simple Brownian diffusion at the same effective diffusion coefficient (about 0.2 $\mu\text{m}^2/\text{s}$). In HS-SPT observations, each glycosphingolipid molecule underwent actin-dependent hop diffusion over the membrane compartment of an average of 110 nm in diameter at an average frequency of once every 25 ms. Surprisingly, this behavior is the same as that of a typical non-raft phospholipid DOPE, qualitatively and quantitatively. These results are at variance with the previous FCS/FRAP studies. In addition, previous investigations concluded that GM₁, labeled either with cholera toxin (CTX) in its headgroup or with Bodipy-FL on the alkyl chain, diffused much more slowly than non-raft phospholipid probes. Here, we found that CTX bound to GM₁ in the plasma membrane diffused as fast as DOPE during the initial 0.5 s after its binding to the membrane, but slowed by a factor of 4 within 10 min after its binding. These results suggest that the slowed diffusion of GM₁ found in previous observations would probably be due to the influences of CTX-induced GM₁ crosslinking or of perturbation by Bodipy-FL on the alkyl chain. These results indicate that in the plasma membrane of non-stimulated cells, glycosphingolipids movements are not slowed by the possible presence of raft, further suggesting that the rafts in the steady-state cells are much smaller than the compartment size (110 nm) and/or short-lived ($<< 25 \text{ ms}$).

1433-Pos Board B277

Role of Membrane Domains in Interferon Receptor Signaling: a Single-molecule Study

Anna Pezzarossa¹, Dirk Schaible², Tobias Meckel¹, Jacob Piehler²,
Thomas Schmidt¹.

¹Leiden University, Leiden, Netherlands, ²University of Osnabruck, Osnabruck, Germany.

Signaling in living cells is largely mediated through multi-protein complexes, and is triggered by recognition of a chemical ligand by membrane receptors at the extracellular side, leading to activation of cytoplasmic effectors. Thus, understanding the dynamical behavior of receptors in the cell membrane is fundamental to understand the processes of cell signaling. The system we are studying, type I interferon (IFN) receptor, is a member of the cytokine family, which plays a key role in early innate and adaptive immune responses upon infection by pathogens. Different members of the type-1 IFN family elicit differential responses although binding to the same receptor. The latter comprises two subunits ifnar1 and ifnar2. Upon ligand binding a ternary complex is formed and signaling pathways activated. Using single-molecule wide-field fluorescence microscopy we follow receptor diffusion in the plasma membrane of living HeLa cells. Receptor subunits are labeled through post-translational labeling with synthetic dyes (e.g. Cy5) coupled to coenzyme-A. Each of the receptor subunit is transfected and expressed, both separately and simultaneously, in HeLa cells, allowing measurements on single component as well as on the ternary complex formed upon IFN binding. Using correlation analysis we obtained information on receptor diffusion constants. Switching between fast and slow motility and vice versa was observed, and interpreted as association/dissociation of ternary complex. Mutants ligand with different affinity toward ifnar1 and ifnar2 are also tested to probe the effect on ternary complex dynamics. Finally we show that membrane nanostructure is possibly involved in the dynamic behavior of the complex. Our data lead to a kinetic model for receptor assembly which may help to obtain a better understanding of trans-membrane signaling.

1434-Pos Board B278

Cortical Cytoskeletal Structures Constrain CD36 Receptor Motion at the Cell Surface to Enhance Aggregation and Signaling

Khuloud Jaqaman¹, Hiro Kuwata², William Trimble², Sergio Grinstein²,
Gaudenz Danuser¹.

¹The Scripps Research Institute, La Jolla, CA, USA, ²Sick Kids, Toronto, ON, Canada.

CD36 is a key receptor in human macrophages, binding to multivalent ligands such as oxidized LDL and malaria-infected red blood cells. To study the dynamics and aggregation kinetics of CD36 receptors in the plasma membrane, we used single-molecule imaging combined with single-particle tracking and mathematical modeling of individual receptor behavior. We immuno-labeled